



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SOME REMARKS ON CONFORMAL REPRESENTATION.*

BY T. H. GRONWALL.

1. In recent investigations of the uniformization of analytic functions, the following proposition† is of fundamental importance:

When the analytic function

$$(1) \quad z = f(x) = \sum_{n=1}^{\infty} a_n x^n$$

effects the conformal representation of the circle $|x| < 1$ on a simple region‡ in the z -plane, the area of this region being less than or equal to A , then, for every $r < 1$, the expressions $|f(x)|$ and $|f'(x)|$ have upper boundaries for $|x| \leq r$ which depend on r and A only, and not on the coefficients in (1).

Koebe gives the inequality

$$|f(x)| < \sqrt{A} \left(\frac{r}{\pi(1-r)^2} + 1 \right) \quad \text{for} \quad |x| \leq r < 1$$

and Courant the corresponding inequality for $f'(x)$

$$|f'(x)| \leq \sqrt{\frac{A}{\pi}} \cdot \frac{1}{1-r^2}.$$

We shall now show that for $|x| \leq r < 1$

$$(2) \quad \begin{aligned} |f(x)| &\leq \sqrt{\frac{A}{\pi}} \cdot \sqrt{\log \frac{1}{1-r^2}}, \\ |f'(x)| &\leq \sqrt{\frac{A}{\pi}} \cdot \frac{1}{1-r^2}, \end{aligned}$$

and that these upper boundaries cannot be replaced by any smaller ones.

Under the assumption made on $f(x)$, the circle $|x| < r$ is conformally represented on a simple region in the z -plane, the area of which is

$$(3) \quad A(r) = \int_0^r d\rho \int_0^{2\pi} |f'(\rho e^{i\theta})|^2 \rho d\theta = \pi \sum_{n=1}^{\infty} n |a_n|^2 r^{2n},$$

and since this region is interior to that corresponding to the circle $|x| < 1$,

* Read before the American Mathematical Society, October 31, 1914.

† Koebe, *Journal für Mathematik*, 138 (1910), p. 222. Courant, *Math. Annalen*, 71 (1912), p. 164.

‡ Simple region = *schlichter Bereich*, a simply connected and nowhere overlapping region.

we have

$$\pi \sum_{n=1}^{\infty} n |a_n|^2 r^{2n} < A$$

for any $r < 1$. A fortiori, the inequality

$$\pi \sum_{n=1}^N n |a_n|^2 r^{2n} < A,$$

subsists, and letting r tend toward unity, we obtain

$$\pi \sum_{n=1}^N n |a_n|^2 \leq A,$$

whence, observing that the terms of the series are positive, we find by increasing N indefinitely

$$(4) \quad \pi \sum_{n=1}^{\infty} n |a_n|^2 \leq A.$$

On the other hand, it is evident that for $|x| \leq r$

$$(5) \quad |f(x)| \leq \sum_{n=1}^{\infty} |a_n| r^n.$$

In the Lagrange inequality

$$\left(\sum_{n=1}^{\infty} u_n v_n \right)^2 \leq \sum_{n=1}^{\infty} u_n^2 \cdot \sum_{n=1}^{\infty} v_n^2,$$

where u_n, v_n are any positive quantities such that the series are convergent, we now make $u_n = \sqrt{n} |a_n|$, $v_n = r^n / \sqrt{n}$ and obtain from (5),

$$|f(x)|^2 \leq \sum_{n=1}^{\infty} n |a_n|^2 \cdot \sum_{n=1}^{\infty} \frac{r^{2n}}{n} = \sum_{n=1}^{\infty} n |a_n|^2 \cdot \log \frac{1}{1-r^2},$$

whence, using (4), we immediately obtain the first inequality (2). To show that the upper boundary obtained is the smallest possible, consider the special function

$$f_0(x) = \sqrt{\frac{A}{\pi}} \cdot \frac{\log \frac{1}{1-r_0 x}}{\sqrt{\log \frac{1}{1-r_0^2}}},$$

where $0 < r_0 < 1$. This function effects the conformal representation of the unit circle on a simple region, since for any two different values of x inside the unit circle the values of $f_0(x)$ are different, and from (3) it is seen at once that the area of this region equals A . Furthermore, for $x = r_0$, we have

$$|f_0(r_0)| = \sqrt{\frac{A}{\pi}} \cdot \sqrt{\log \frac{1}{1-r_0^2}},$$

so that the upper boundary (2) is actually reached for the particular value $r = r_0$.

To prove the second inequality (2), we observe that for $|x| \leq r$

$$|f'(x)| \leq \sum_{n=1}^{\infty} n |a_n| r^{n-1};$$

applying Lagrange's inequality with $u_n = \sqrt{n} |a_n|$, $v_n = \sqrt{n} r^{n-1}$, we find

$$|f'(x)|^2 \leq \sum_{n=1}^{\infty} n |a_n|^2 \cdot \sum_{n=1}^{\infty} n r^{2n-2} = \sum_{n=1}^{\infty} n |a_n|^2 \cdot \frac{1}{(1-r^2)^2},$$

and combining with (4), the desired result is obtained. Using the comparison function

$$f_0(x) = (1 - r_0^2) \sqrt{\frac{A}{\pi}} \cdot \frac{x}{1 - r_0 x}$$

it is seen as before that the upper boundary cannot be replaced by any smaller one.

2. In connection with his exposition of Koebe's distortion theorem, Fricke* proves the following proposition:

When

$$(6) \quad f(x) = \frac{1}{x} + \sum_{n=1}^{\infty} a_n x^n,$$

where the power series has no constant term and converges for $|x| < 1$, and $f(x)$ does not take the same value for two different values of x inside the unit circle, then the maximum $M(r)$ of $|f(x)|$ for $|x| = r$ satisfies the inequality

$$M(r) < \frac{2 + 2\sqrt{2}}{r}, \quad 0 < r < 1.$$

We shall now show that this inequality may be replaced by

$$(7) \quad M(r) < \frac{9}{4} \cdot \frac{1}{r}.$$

Since $f(x)$ takes each of its values but once in the unit circle, $z = f(x)$ effects the conformal representation of the unit circle on a simple region in the z -plane, containing the point $z = \infty$ (corresponding to $x = 0$) in its interior; to the circumference $|x| = r$ ($0 < r < 1$) there corresponds a closed curve C_r , all the points of which are at finite distance, and if $r > r_0$, the curve C_r lies entirely inside the curve C_{r_0} . Consequently

* R. Fricke and F. Klein, Vorlesungen über die Theorie der automorphen Functionen, vol. 2 (Leipzig, Teubner, 1912), pp. 497-498.

$M(r) < M(r_0)$, and

$$rM(r) < rM(r_0) < M(r_0), \quad 0 < r_0 < r < 1.$$

But since $xf(x)$ is holomorphic for $|x| < 1$, we have for $r \leq r_0$

$$rM(r) = \max_{|x|=r} |xf(x)| \leq \max_{|x|=r_0} |xf(x)| = r_0 M(r_0) < M(r_0),$$

and this, together with the preceding inequality, gives

$$(8) \quad rM(r) < M(r_0), \quad 0 \leq r < 1.$$

We shall now establish an upper boundary for $M(r_0)$, and then determine r_0 so as to make this boundary as small as possible. The area inside the curve C_r is readily found to be

$$\pi \left(1 - \sum_{n=1}^{\infty} n |a_n|^2 r^{2n} \right),$$

and this area being obviously positive for every $r < 1$, we find, letting r tend toward unity in the same way as when proving (4), that

$$(9) \quad \sum_{n=1}^{\infty} n |a_n|^2 \leq 1.$$

Now

$$M(r) \leq \frac{1}{r} + \sum_{n=1}^{\infty} |a_n| r^n,$$

and applying the Lagrange inequality and the relation (9) to the right hand series in the same way as before, we see that

$$M(r) \leq \frac{1}{r} + \sqrt{\log \frac{1}{1-r^2}}.$$

In the interval $0 \leq r \leq 1$, the last expression has only one minimum = 2.242 ... for $r = 0.74$...; taking the latter value for r_0 , it follows from (8) that

$$rM(r) < 2.242 \dots < \frac{9}{4} \quad \text{for} \quad 0 < r < 1,$$

which proves our statement (6).

The particular function

$$(10) \quad f(x) = \frac{1}{x} + e^{2\alpha i} x,$$

where α is real, effects a conformal representation of the unit circle on that part of the z -plane which lies outside of the rectilinear cut from

$z = -2e^{ai}$ to $z = 2e^{ai}$. For this function we have

$$M(r) = \frac{1}{r} + r \leq \frac{2}{r} \quad \text{for} \quad 0 < r \leq 1,$$

so that the constant in (7) cannot be replaced by any value less than 2. It is probable that, for every function (6) satisfying the conditions stated, $M(r) \leq \frac{1}{r} + r$, the equality sign holding only when $f(x)$ has the form (10).

The proof of this statement seems very difficult; the case, however, when all a_n are real and not negative, may be disposed of as follows. From the assumptions concerning $f(x)$, it is seen that $f'(x) \neq 0$ inside the unit circle, and consequently

$$f'(r) = -\frac{1}{r^2} + \sum_{n=1}^{\infty} n a_n r^{n-1} < 0 \quad \text{for} \quad 0 < r < 1,$$

since otherwise $f'(x) = 0$ would have a positive root $\leq r$. Consequently, letting r tend toward unity in the same way as before, we find

$$\sum_{n=1}^{\infty} n a_n \leq 1,$$

or

$$a_1 \leq 1 - \sum_{n=2}^{\infty} n a_n,$$

whence

$$M(r) = \frac{1}{r} + \sum_{n=1}^{\infty} a_n r^n \leq \frac{1}{r} + r - \sum_{n=2}^{\infty} a_n (nr - r^n) \leq \frac{1}{r} + r,$$

the equality sign holding only for $a_1 = 1$ and $a_n = 0$, ($n > 1$).

PRINCETON UNIVERSITY.